



Estimated Age-Standardized Prevalence of Daily Smoking and Annualized Rate of Change, 1980-2012See the Supplement for full list of developing and developed countries, as defined by the Global Burden of Disease Study. For 6 countries without data on tobacco smoking prevalence (Afghanistan, Angola, Central African Republic, Guineau-Bissau, Somalia, Turkmenistan), estimates were derived based on tobacco consumption and trends in prevalence in neighboring countries. The Supplement has more details on these methods. A, Global (187 countries) age-standardized prevalence estimates by sex and separately for developed countries (50 countries) and developing countries (137 countries) from 1980 to 2012. B, Annualized rate of change in age-standardized prevalence by sex from 1980 to 2012. Annualized rates of change are plotted at the beginning of the time interval they correspond to (eg, the rate of change shown for 2011 is the rate of change between 2011 and 2012). Shaded areas designate 95% uncertainty intervals. Interactive data display.


Source: Smoking prevalence and cigarette consumption in 187 countries, 1980-2012. JAMA, 2014, 311: 183-192.



Source: Smoking prevalence and cigarette consumption in 187 countries, 1980-2012. JAMA, 2014, 311: 183-192.




A Probable mental illness (any diagnosis or PHQ2>2 or K6>12)



Source: Trends in smoking among adults with mental illness and association between mental health treatment and smoking cessation. *JAMA*, 2014, 311: 172-182.

INTENTION TO QUIT SMOKING



Source: Eurobarometer 385, http://ec.europa.eu/public_opinion/index_en.htm